

$SU(3)$ mass-splittings of heavy-baryons in QCD

R.M. Albuquerque^{*a}, S. Narison^{†b}, M. Nielsen^a,

^aInstituto de Física, Universidade de São Paulo, C.P. 66318, 05389-970 São Paulo, SP, Brazil.

^bLaboratoire de Physique Théorique et Astroparticules, CNRS-IN2P3 & Université de Montpellier II, Case 070, Place Eugène Bataillon, 34095 - Montpellier Cedex 05, France.

We extract directly (for the first time) the charmed ($C = 1$) and bottom ($B = -1$) heavy-baryons (spin 1/2 and 3/2) mass-splittings due to $SU(3)$ breaking using double ratios of QCD spectral sum rules (QSSR) in full QCD, which are less sensitive to the exact value and definition of the heavy quark mass, to the perturbative radiative corrections and to the QCD continuum contributions than the simple ratios commonly used for determining the heavy baryon masses. Noticing that most of the mass-splittings are mainly controlled by the ratio $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$ of the condensate, we extract this ratio, by allowing 1σ deviation from the observed masses of the $\Xi_{c,b}$ and of the Ω_c . We obtain: $\kappa = 0.74(3)$, which improves the existing estimates: $\kappa = 0.70(10)$ from light hadrons. Using this value, we deduce $M_{\Omega_b} = 6078.5(27.4)$ MeV which agrees with the recent CDF data but disagrees by 2.4σ with the one from D0. Predictions of the Ξ'_Q and of the spectra of spin 3/2 baryons containing one or two strange quark are given in Table 2. Predictions of the hyperfine splittings $\Omega_Q^* - \Omega_Q$ and $\Xi_Q^* - \Xi_Q$ are also given in Table 3. Starting for a general choice of the interpolating currents for the spin 1/2 baryons, our analysis favours the optimal value of the mixing angle $b \simeq (-1/5 \sim 0)$ found from light and non-strange heavy baryons.

1. Introduction

QSSR [1,2] à la SVZ [3] has been used earlier in full QCD [4,5,6] and in HQET [7] for understanding heavy baryons [charmed (cqq), bottom (bqq), double charm (ccq), double bottom (bbq) and (bcq)] masses. Recent observations at Tevatron of families of b -baryons [8,9] and of the Ω_c^* baryon by Babar and Belle [10] have stimulated different recent theoretical activities for understanding their nature [11,12,13,14,15,16,17,18,19]. QSSR results are in quite good agreement with recent experimental findings but with relatively large uncertainties. The inaccuracy of these results is mainly due to the value of the heavy quark mass and of its ambiguous definition when working to lowest order (LO) in the radiative α_s corrections in full QCD and HQET¹, where the heavy quark mass is the main driving term in the QCD expression of the baryon two-point correlator used in the QSSR analysis. Another source of uncertainty is the effect of the QCD continuum which parametrizes the higher baryon masses contributions to the spectral function and the *ad hoc* choices of interpolating baryon currents used in different literatures. In this paper, we shall concentrate on the analysis of the heavy baryons mass-splittings due to $SU(3)$ breaking using double ratios (DR) of QCD spectral sum rules (QSSR), which are less sensitive to the exact value and definition of the heavy quark mass and to the QCD continuum contributions than the simple ratios used in the literature to determine the absolute value of heavy baryon masses.

In this letter, we extend the previous analysis in [4,5] by including the new $SU(3)$ breaking terms: m_s and the ratio of the condensate $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$.

For the spin 1/2 baryons, and following Ref. [4], we

work with the lowest dimension general currents:

$$\begin{aligned} \eta_{\Xi_Q} &= \epsilon_{abc} [(q_a^T C \gamma_5 s_b) + b(q_a^T C s_b) \gamma_5] Q_c, \\ \eta_{\Lambda_Q} &= \eta_{\Xi_Q} \quad (s \rightarrow q), \\ \eta_{\Omega_Q} &= \epsilon_{abc} [(s_a^T C \gamma_5 Q_b) + b(s_a^T C Q_b) \gamma_5] s_c, \\ \eta_{\Sigma_Q} &= \eta_{\Omega_Q} \quad (s \rightarrow q), \\ \eta_{\Xi'_Q} &= \frac{1}{\sqrt{2}} \epsilon_{abc} \left[(s_a^T C \gamma_5 Q_b) q_c + (q_a^T C \gamma_5 Q_b) s_c \right. \\ &\quad \left. + b((s_a^T C Q_b) \gamma_5 q_c + (q_a^T C Q_b) \gamma_5 s_c) \right], \end{aligned} \quad (1)$$

where we use standard notations; b is *a priori* an arbitrary mixing parameter. Its value has been found to be:

$$b = -1/5, \quad (2)$$

in the case of light baryons [20] and in the range [4,5,6]:

$$-0.5 \leq b \leq 0.5, \quad (3)$$

for non-strange heavy baryons, which do not favour the Ioffe choice $b = -1$ [21]. The corresponding two-point correlator reads:

$$\begin{aligned} S(q) &= i \int d^4x e^{iqx} \langle 0 | T \bar{\eta}_Q(x) \eta_Q(0) | 0 \rangle \\ &\equiv \hat{q} F_1 + F_2, \end{aligned} \quad (4)$$

where F_1 and F_2 are two invariant functions.

For the spin 3/2 baryons, we follow Ref. [5] and work with the interpolating currents:

$$\begin{aligned} \eta_{\Xi_Q^*}^\mu &= \sqrt{\frac{2}{3}} [(q^T C \gamma_\mu Q) s + (s^T C \gamma_\mu Q) q + (q^T C \gamma_\mu s) Q] \\ \eta_{\Omega_Q^*}^\mu &= \frac{1}{\sqrt{2}} \eta_{\Xi_Q^*}^\mu \quad (q \rightarrow s) \\ \eta_{\Sigma_Q^*}^\mu &= \frac{1}{\sqrt{2}} \eta_{\Xi_Q^*}^\mu \quad (s \rightarrow q), \end{aligned} \quad (5)$$

where an anti-symmetrization over colour indices is understood. The normalization in Eq. (5) is chosen in such

^{*}E-mail addresses: rma@if.usp.br (R.M. Albuquerque), snarison@yahoo.fr (S. Narison), mnielsen@if.usp.br (M. Nielsen).

[†]Corresponding author.

¹Radiative corrections to the heavy baryon correlators have been evaluated in [16] but for a particular choice of the interpolating currents.

a way that in all cases one gets the same perturbative contribution. The corresponding two-point correlator reads:

$$\begin{aligned} S^{\mu\nu}(q) &= i \int d^4x e^{iqx} \langle 0 | T \bar{\eta}_Q^\mu(x) \eta_Q^\nu(0) | 0 \rangle \\ &\equiv g^{\mu\nu} (\hat{q}F_1 + F_2) + \dots \end{aligned} \quad (6)$$

In the following, the contribution of the heavy quark condensate $m_Q \langle \bar{Q}Q \rangle$ will not appear [5] as it is cancelled by a part of the gluon condensate contribution due to the heavy quark relation $m_Q \langle \bar{Q}Q \rangle + (1/12\pi) \langle \alpha_s G^2 \rangle \simeq 0$ [3,1]. Again due to this relation, the one due to $m_s \langle \bar{Q}Q \rangle$ is numerically negligible because of the extra $(1/12\pi)$ loop and $1/M_Q$ factors compared to the one due to $m_s \langle \bar{q}q \rangle$ and due to the four-quark condensate contributions. The same loop factor also numerically suppresses the contributions of $\langle \alpha_s G^2 \rangle$ and $m_s \langle \alpha_s G^2 \rangle$ compared to the other ones. These negligible $SU(3)$ breaking contributions will not be considered in the following.

2. The spin 1/2 two-point correlator in QCD

The $\Lambda_Q(Qqq)$ and $\Xi_Q(Qsq)$ baryons

The expression for Λ_Q has been (first) obtained in the chiral limit $m_q = 0$ in [5], and the one of Ξ_Q including $SU(3)$ breaking in [14]. One can notice that due to the expression of the current the m_s corrections vanish to leading order in α_s for the perturbative term, while the $D = 6$ condensates for the $SU(2)$ case of [5] needs the following replacement in the $SU(3)$ case:

$$\rho \langle \bar{q}q \rangle^2 \rightarrow \rho \langle \bar{q}q \rangle \langle \bar{s}s \rangle, \quad (7)$$

where $\rho = 2 \sim 3$ indicates the violation of the four-quark vacuum saturation [22,1,30]. The additional $SU(3)$ breaking corrections for the Ξ_Q are [14]:

- F_1 :

$$\begin{aligned} \text{Im} F_1^{m_s} |_{\bar{s}s} &= -\frac{m_s}{2^4 \pi} (1-x^2) \left[(1-b^2) \langle \bar{q}q \rangle - \frac{(1+b^2)}{2} \langle \bar{s}s \rangle \right], \\ F_1^{m_s} |_{mix} &= \frac{m_s}{2^5 \pi^2} \frac{1}{m_Q^2 - q^2} \left\{ \langle \bar{s}Gs \rangle \frac{(1+b^2)}{6} + \langle \bar{q}Gq \rangle (1-b^2) \right\}, \end{aligned} \quad (8)$$

- F_2 :

$$\begin{aligned} \text{Im} F_2^{m_s} |_{\bar{s}s} &= -\frac{m_s m_Q}{2^3 \pi} (1-x) \left[(1+b^2) \langle \bar{q}q \rangle - \frac{(1-b^2)}{2} \langle \bar{s}s \rangle \right], \\ F_2^{m_s} |_{mix} &= \frac{m_s m_Q}{2^5 \pi^2} \frac{1}{m_Q^2 - q^2} \left\{ \langle \bar{s}Gs \rangle \frac{(1-b^2)}{6} + \langle \bar{q}Gq \rangle (1+b^2) \right\}, \end{aligned} \quad (9)$$

where $x \equiv m_Q^2/s$ and $\langle \bar{s}Gs \rangle \equiv g \langle \bar{s} \sigma_{\mu\nu} \lambda_a / 2 G_a^{\mu\nu} s \rangle$.

The $\Sigma_Q(Qqq)$ and $\Omega_Q(Qss)$ baryons

The expression for Σ_Q has been (first) obtained in [4]. The additional $SU(3)$ breaking terms for the Ω_Q are:

- F_1 :

$$\begin{aligned} \text{Im} F_1^{m_s} |_{pert} &= \frac{3m_s m_Q^3}{2^8 \pi^3} (1-b^2) \times \left[\frac{2}{x} + 3 - 6x + x^2 + 6 \ln x \right], \\ \text{Im} F_1^{m_s} |_{\bar{s}s} &= \frac{3m_s \langle \bar{s}s \rangle}{2^6 \pi} (1+b)^2 (1-x^2), \\ F_1^{m_s} |_{mix} &= -\frac{m_s \langle \bar{s}Gs \rangle}{2^7 3 \pi^2} \left[\frac{1}{m_Q^2 - q^2} (7 + 22b + 7b^2) - 6(1+b)^2 \int_0^1 \frac{d\alpha (1-\alpha)}{m_Q^2 - (1-\alpha)q^2} \right], \\ F_1^{m_s} |_{D=6} &= -\frac{m_s m_Q \rho \langle \bar{s}s \rangle^2 (1-b^2)}{8(m_Q^2 - q^2)^2}. \end{aligned} \quad (10)$$

- F_2 :

$$\begin{aligned} \text{Im} F_2^{m_s} |_{pert} &= \frac{3m_s m_Q^4}{2^8 \pi^3} (1-b^2) \times \left(\frac{1}{x^2} - \frac{6}{x} + 3 + 2x - 6 \ln x \right), \\ \text{Im} F_2^{m_s} |_{\bar{s}s} &= -\frac{3m_s m_Q \langle \bar{s}s \rangle}{2^5 \pi} (3 + 2b + 3b^2) (1-x), \\ F_2^{m_s} |_{mix} &= \frac{m_s m_Q \langle \bar{s}Gs \rangle}{2^7 3 \pi^2} \times \left[\frac{1}{m_Q^2 - q^2} (25 + 22b + 25b^2) - 3(5 + 6b + 5b^2) \times \int_0^1 \frac{d\alpha}{m_Q^2 - (1-\alpha)q^2} \right], \\ F_2^{m_s} |_{D=6} &= -\frac{m_s \rho \langle \bar{s}s \rangle^2 (1-b^2)}{8(m_Q^2 - q^2)} \left[1 + \frac{m_Q^2}{m_Q^2 - q^2} \right], \end{aligned} \quad (11)$$

The $\Sigma_Q(Qqq)$ and $\Xi'_Q(Qsq)$ baryons

The expression for the Ξ'_Q tends to the one of the Σ_Q in the chiral limit $m_{q,s} \rightarrow 0$ and is very similar with one of the Ω_Q . The $SU(3)$ breaking corrections read:

F_1 :

$$\begin{aligned} \text{Im} F_1^{m_s} |_{pert} &= \frac{3m_s m_Q^3}{2^9 \pi^3} (1-b^2) \times \left[\frac{2}{x} + 3 - 6x + x^2 + 6 \ln x \right], \\ \text{Im} F_1 |_{\bar{s}s} &= -\frac{3m_Q (\langle \bar{s}s \rangle + \langle \bar{q}q \rangle)}{2^6 \pi} (1-b^2) (1-x)^2, \\ \text{Im} F_1^{m_s} |_{\bar{s}s} &= \frac{m_s}{2^7 \pi} (1-x^2) \left[-2(1-b)^2 \langle \bar{q}q \rangle + (5 + 2b + 5b^2) \langle \bar{s}s \rangle \right], \end{aligned}$$

$$\begin{aligned}
\text{Im } F_1|_{mix} &= \frac{(\langle \bar{s}Gs \rangle + \langle \bar{q}Gq \rangle)}{m_Q 2^8 \pi} (1-b^2) (13x^2 - 6x), \\
F_1^{m_s}|_{mix} &= -\frac{m_s}{2^8 3\pi^2} \left[\frac{1}{m_Q^2 - q^2} \left[(13 + 10b + 13b^2) \times \right. \right. \\
&\quad \left. \langle \bar{s}Gs \rangle - 6(1-b)^2 \langle \bar{q}Gq \rangle \right] + \\
&\quad \left[3(1-b)^2 \langle \bar{q}Gq \rangle \right. \\
&\quad \left. - 3(3+2b+3b^2) \langle \bar{s}Gs \rangle \right] \times \\
&\quad \left. \int_0^1 d\alpha \frac{1-\alpha}{m_Q^2 - (1-\alpha)q^2} \right], \\
F_1|_{D=6} &= \frac{\rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{24} (1-b)^2 \frac{1}{m_Q^2 - q^2}, \\
F_1^{m_s}|_{D=6} &= -\frac{m_s m_Q \rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{16} \frac{1-b^2}{(m_Q^2 - q^2)^2} \quad (12)
\end{aligned}$$

F_2 :

$$\begin{aligned}
\text{Im } F_2^{m_s}|_{pert} &= \frac{3m_s m_Q^4}{2^9 \pi^3} (1-b^2) \times \\
&\quad \left[\frac{1}{x^2} - \frac{6}{x} + 3 + 2x - 6 \ln x \right], \\
\text{Im } F_2|_{\bar{s}s} &= -\frac{3m_Q^2 (\langle \bar{s}s \rangle + \langle \bar{q}q \rangle)}{2^6 \pi} \times \\
&\quad (1-b^2)x \left(1 - \frac{1}{x} \right)^2, \\
\text{Im } F_2^{m_s}|_{\bar{s}s} &= \frac{m_s m_Q}{2^6 \pi} (1-x) \left[(1-b)^2 \langle \bar{s}s \rangle - \right. \\
&\quad \left. 2(5+2b+5b^2) \langle \bar{q}q \rangle \right], \\
\text{Im } F_2|_{mix} &= \frac{(\langle \bar{s}Gs \rangle + \langle \bar{q}Gq \rangle)}{2^8 \pi} (1-b^2) (6+x), \\
F_2^{m_s}|_{mix} &= -\frac{m_s m_Q}{2^8 3\pi^2} \left[\frac{1}{m_Q^2 - q^2} \left[5(1-b)^2 \langle \bar{s}Gs \rangle \right. \right. \\
&\quad \left. - 6(5+2b+5b^2) \langle \bar{q}Gq \rangle \right] + \\
&\quad \left[-3(1-b)^2 \langle \bar{s}Gs \rangle + \right. \\
&\quad \left. 6(3+2b+3b^2) \times \right. \\
&\quad \left. \langle \bar{q}Gq \rangle \right] \int_0^1 \frac{d\alpha}{m_Q^2 - (1-\alpha)q^2}, \\
F_2|_{D=6} &= \frac{m_Q \rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{24} (5+2b+5b^2) \frac{1}{m_Q^2 - q^2}, \\
F_2^{m_s}|_{D=6} &= -\frac{m_s \rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{16} \frac{1-b^2}{m_Q^2 - q^2} \times \\
&\quad \left[1 + \frac{m_Q^2}{m_Q^2 - q^2} \right]. \quad (13)
\end{aligned}$$

We have checked the existing results in [4] obtained in the chiral limit and all our previous results agree with these ones.

3. The spin 3/2 two-point correlator in QCD

The QCD expression of the two-point correlator for the $\Sigma_Q^*(Qqq)$ has been (first) obtained in the chiral limit

$m_{u,d} = 0$, to LO in α_s and up to the contributions of the $D = 6$ condensates in [4]. In this letter, we extend the previous analysis by including the new $SU(3)$ breaking m_s correction terms and consider the $SU(3)$ breaking of the ratio of quark condensates $\langle \bar{s}s \rangle \neq \langle \bar{q}q \rangle$ like we did for the spin 1/2 case.

The $\Sigma_Q^*(Qqq)$ and $\Xi_Q^*(Qsq)$ baryons

The additionnal terms and replacement due to $SU(3)$ breaking for the Ξ_Q^* compared with the one of the $\Sigma_Q^*(Qqq)$ in [5] are:

- F_1 :

$$\begin{aligned}
\text{Im } F_1^{m_s}|_{pert} &= \frac{m_s m_Q^3}{48 \pi^3} \left[\frac{2}{x} + 3 - 6x + x^2 + 6 \ln x \right], \\
\text{Im } F_1|_{\bar{s}s} &= -\frac{m_Q}{6\pi} \left[\langle \bar{q}q \rangle + \langle \bar{s}s \rangle \right] (1-x)^2, \\
\text{Im } F_1^{m_s}|_{\bar{s}s} &= -\frac{m_s}{12\pi} \left[2(1-x^2) \langle \bar{q}q \rangle - (1-x^3) \langle \bar{s}s \rangle \right], \\
\text{Im } F_1|_{mix} &= \frac{7M_0^2}{3^2 2^3 \pi} \left[\langle \bar{q}q \rangle + \langle \bar{s}s \rangle \right] \frac{x^2}{m_Q}, \\
F_1^{m_s}|_{mix} &= \frac{m_s M_0^2}{144 \pi^2} \left[\frac{12 \langle \bar{q}q \rangle - 9 \langle \bar{s}s \rangle}{m_Q^2 - q^2} + \right. \\
&\quad \left. 2 \int_0^1 \frac{d\alpha (1-\alpha)}{m_Q^2 - (1-\alpha)q^2} \times \right. \\
&\quad \left. \left[(1-3\alpha) \langle \bar{s}s \rangle + \langle \bar{q}q \rangle \right] \right], \\
F_1|_{D=6} &= \frac{4}{9} \frac{\rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{m_Q^2 - q^2}, \\
F_1^{m_s}|_{D=6} &= -\frac{2}{9} m_Q m_s \frac{\rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{(m_Q^2 - q^2)^2}. \quad (14)
\end{aligned}$$

- F_2 :

$$\begin{aligned}
\text{Im } F_2^{m_s}|_{pert} &= \frac{m_s m_Q^4}{192 \pi^3} \left[\frac{3}{x^2} - \frac{16}{x} + 12 + x^2 - 12 \ln x \right], \\
\text{Im } F_2|_{\bar{s}s} &= -\frac{m_Q^2}{18\pi} \left[\langle \bar{q}q \rangle + \langle \bar{s}s \rangle \right] \left(\frac{2}{x} - 3 + x^2 \right), \\
\text{Im } F_2^{m_s}|_{\bar{s}s} &= -\frac{m_s m_Q}{12\pi} (1-x) \left[6 \langle \bar{q}q \rangle - (1+x) \langle \bar{s}s \rangle \right], \\
\text{Im } F_2|_{mix} &= \frac{M_0^2}{18\pi} \left[\langle \bar{q}q \rangle + \langle \bar{s}s \rangle \right] \left(1 + \frac{3}{4} x^2 \right), \\
F_2^{m_s}|_{mix} &= \frac{m_s m_Q M_0^2}{72 \pi^2} \left[3 \frac{(3 \langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{m_Q^2 - q^2} + \right. \\
&\quad \left. \langle \bar{q}q \rangle \int_0^1 \frac{d\alpha}{m_Q^2 - (1-\alpha)q^2} \right], \\
F_2|_{D=6} &= \frac{2}{3} \frac{m_Q \rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{m_Q^2 - q^2}, \\
F_2^{m_s}|_{D=6} &= -\frac{2}{9} \frac{m_s m_Q^2 \rho \langle \bar{s}s \rangle \langle \bar{q}q \rangle}{(m_Q^2 - q^2)^2}, \quad (15)
\end{aligned}$$

where $x \equiv m_Q^2/s$ and $\langle \bar{s}Gs \rangle \equiv g \langle \bar{s} \sigma_{\mu\nu} \lambda_a / 2 G_a^{\mu\nu} s \rangle \equiv M_0^2 \langle \bar{s}s \rangle$.

The $\Omega_Q^*(Qss)$ baryons

Compared with the expression of the $\Sigma_Q^*(Qqq)$ in [5], the additionnal $SU(3)$ breaking terms for the Ω_Q^* are:

- F_1 :

$$\begin{aligned} \text{Im}F_1^{m_s}|_{\text{pert}} &= \frac{m_s m_Q^3}{24\pi^3} \left[\frac{2}{x} + 3 - 6x + x^2 + 6 \ln x \right], \\ \text{Im}F_1^{m_s}|_{\bar{s}s} &= -\frac{m_s \langle \bar{s}s \rangle}{6\pi} (1 - 2x^2 + x^3), \\ F_1^{m_s}|_{\text{mix}} &= \frac{m_s M_0^2 \langle \bar{s}s \rangle}{72\pi^2} \left[\frac{3}{m_Q^2 - q^2} + \right. \\ &\quad \left. 2 \int_0^1 \frac{d\alpha (1-\alpha)(2-3\alpha)}{m_Q^2 - (1-\alpha)q^2} \right], \\ F_1^{m_s}|_{D=6} &= -\frac{4 m_Q m_s \rho \langle \bar{s}s \rangle^2}{9 (m_Q^2 - q^2)^2}. \end{aligned} \quad (16)$$

- F_2 :

$$\begin{aligned} \text{Im}F_2^{m_s}|_{\text{pert}} &= \frac{m_s m_Q^4}{96\pi^3} \\ &\quad \left[\frac{3}{x^2} - \frac{16}{x} + 12 + x^2 - 12 \ln x \right], \\ \text{Im}F_2^{m_s}|_{\bar{s}s} &= -\frac{m_s m_Q \langle \bar{s}s \rangle}{6\pi} \left[5 - 6x + x^2 \right], \\ F_2^{m_s}|_{\text{mix}} &= \frac{m_s m_Q M_0^2 \langle \bar{s}s \rangle}{36\pi^2} \left[\frac{6}{m_Q^2 - q^2} + \right. \\ &\quad \left. \int_0^1 \frac{d\alpha}{m_Q^2 - (1-\alpha)q^2} \right], \\ F_2^{m_s}|_{D=6} &= -\frac{4 m_s m_Q^2 \rho \langle \bar{s}s \rangle^2}{9 (m_Q^2 - q^2)^2}. \end{aligned} \quad (17)$$

We have checked the existing results in [5] obtained in the chiral and $SU(2)$ limits and agree with these ones.

4. Form of the sum rules and QCD inputs

We parametrize the spectral function using the standard duality ansatz: “one resonance” + “QCD continuum”. The QCD continuum starts from a threshold t_c and comes from the discontinuity of the QCD diagrams. Transferring its contribution to the QCD side of the sum rule, one obtains the finite energy Laplace/Borel sum rules:

$$\begin{aligned} |\lambda_{B_q^{(*)}}|^2 M_{B_q^{(*)}} e^{-M_{B_q^{(*)}}^2 \tau} &= \int_{t_q}^{t_c} ds e^{-s\tau} \frac{1}{\pi} \text{Im}F_2(s), \\ |\lambda_{B_q^*}|^2 e^{-M_{B_q^*}^2 \tau} &= \int_{t_q}^{t_c} ds e^{-s\tau} \frac{1}{\pi} \text{Im}F_1(s), \end{aligned} \quad (18)$$

where $\tau \equiv 1/M^2$ is the sum rule variable and $\lambda_{B_q^{(*)}}$ and $M_{B_q^{(*)}}$ are the heavy baryon residue and mass. Notice that, within our choice of the interpolating currents, we may have a contamination due to the negative parity states contribution, which we can quantify from the ratio of sum rules associated to $(M_Q F_1 - F_2)$ and $(M_Q F_1 + F_2)$ [7]. We shall check (a posteriori) that this ratio is negligible at the stability regions. On the

other, the observed negative parity states are systematically heavier than the positive ones such that they can be legitimately included into the QCD continuum contribution. Consistently, we also take into account the $SU(3)$ breaking at the quark and continuum threshold:

$$\begin{aligned} \sqrt{t_q}|_{SU(3)} &\simeq (\sqrt{t_q}|_{SU(2)} \equiv m_Q) + \bar{m}_{q_1} + \bar{m}_{q_2}, \\ \sqrt{t_c}|_{SU(3)} &\simeq (\sqrt{t_c}|_{SU(2)} \equiv \sqrt{t_c}) + \bar{m}_{q_1} + \bar{m}_{q_2}, \end{aligned} \quad (19)$$

where $q_{1,2} \equiv q$ or/and s depending on the channel. $\bar{m}_{q_i}$ are the running light quark masses. m_Q is the heavy quark mass, which we shall take in the range covered by the running and on-shell mass (see Table 1) because of its ambiguous definition when working to lowest order of perturbative QCD. One can estimate the baryon masses from the following ratios:

$$\begin{aligned} \mathcal{R}_i^q &= \frac{\int_{t_q}^{t_c} ds s e^{-s\tau} \text{Im}F_i(s)}{\int_{t_q}^{t_c} ds e^{-s\tau} \text{Im}F_i(s)}, \quad i = 1, 2, \\ \mathcal{R}_{21}^q &= \frac{\int_{t_q}^{t_c} ds e^{-s\tau} \text{Im}F_2(s)}{\int_{t_q}^{t_c} ds e^{-s\tau} \text{Im}F_1(s)}, \end{aligned} \quad (20)$$

where at the τ -stability point :

$$M_{B_q^{(*)}} \simeq \sqrt{\mathcal{R}_i^q} \simeq \mathcal{R}_{21}^q. \quad (21)$$

These quantities have been used in the literature for getting the baryon masses and lead to a typical uncertainty of 15-20% [4,5,6]². In order to circumvent these problems, we work with the double ratio of sum rules (DR)[23]:

$$r_i^{sd} \equiv \sqrt{\frac{\mathcal{R}_i^s}{\mathcal{R}_i^d}}, \quad r_{21}^{sd} \equiv \frac{\mathcal{R}_{21}^s}{\mathcal{R}_{21}^d}. \quad (22)$$

which take directly into account the $SU(3)$ breaking effects. These quantities are obviously less sensitive to the choice of the heavy quark masses, to the perturbative radiative corrections and to the value of the continuum threshold than the simple ratios $\mathcal{R}_i$ and $\mathcal{R}_{21}$ ^{2,3}. Analogous DR quantities have been used successfully (for the first time) in [23] for studying the mass ratio of the $0^{++}/0^{-+}$ and $1^{++}/1^{--}$ B-mesons, in [24] for extracting f_{B_s}/f_B , in [25] for estimating the $D \rightarrow K/D \rightarrow \pi$ semi-leptonic form factors and in [26] for extracting the strange quark mass from the $e^+e^- \rightarrow I = 1, 0$ data. For the numerical analysis we shall introduce the RGI quantities $\hat{\mu}$ and $\hat{m}_q$ [27]:

$$\begin{aligned} \bar{m}_q(\tau) &= \frac{\hat{m}_q}{(-\log \sqrt{\tau} \Lambda)^{2/\beta_1}} \\ \langle \bar{q}q \rangle(\tau) &= -\hat{\mu}_q^3 (-\log \sqrt{\tau} \Lambda)^{2/\beta_1} \\ \langle \bar{q}Gq \rangle(\tau) &= -\hat{\mu}_q^3 (-\log \sqrt{\tau} \Lambda)^{1/-3\beta_1} M_0^2, \end{aligned} \quad (23)$$

²More accurate results quoted in the recent QSSR literature [14, 15] do not take into account the uncertainties due to the heavy quark mass definitions and to the arbitrary choice of the baryonic interpolating currents.

³One may also work with the double ratio of moments $\mathcal{M}_n$ based on different derivatives at $q^2 = 0$ [23]. However, in this case the OPE is expressed as an expansion in $1/m_Q$, which for a LO expression of the QCD correlator is more affected by the definition of the heavy quark mass to be used.

where $\beta_1 = -(1/2)(11 - 2n/3)$ is the first coefficient of the β function for n flavours. We have used the quark mass and condensate anomalous dimensions reviewed in [1]. We shall use the QCD parameters in Table 1. At the scale where we shall work, and using the parameters in the table, we deduce:

$$\rho = 2.1 \pm 0.2, \quad (24)$$

which controls the deviation from the factorization of the four-quark condensates. We shall not include the $1/q^2$ term discussed in [28,29], which is consistent with the LO approximation used here as the latter has been motivated for a phenomenological parametrization of the larger order terms of the QCD series.

Table 1

QCD input parameters. For the heavy quark masses, we use the range spanned by the running $\overline{MS}$ mass $\overline{m}_Q(M_Q)$ and the on-shell mass from QSSR compiled in page 602,603 of the book in [1].

Parameters	Values	Ref.
Λ	(353 ± 15) MeV	[30,9]
$\hat{m}_d$	(6.1 ± 0.5) MeV	[31,1,9]
$\hat{m}_s$	(114.5 ± 20.8) MeV	[31,1,9]
$\hat{\mu}_d$	(263 ± 7) MeV	[31,1]
$\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$	(0.7 ± 0.1)	[31,1,20]
M_0^2	(0.8 ± 0.1) GeV ²	[20,21,23]
$\langle \alpha_s G^2 \rangle$	$(6.8 \pm 1.3) \times 10^{-2}$ GeV ⁴	[30,22,32,33,34,35,2]
$\rho \alpha_s \langle \bar{d}d \rangle^2$	$(4.5 \pm 0.3) \times 10^{-4}$ GeV ⁶	[30,22]
m_c	$(1.18 \sim 1.47)$ GeV	[1,31,36,9]
m_b	$(4.18 \sim 4.72)$ GeV	[1,31,36,9]

5. $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$ from the spin 1/2 baryons

As a preliminary step of the analysis, we check the different results obtained in full QCD and in the chiral limit [4,5]:

$$\begin{aligned} M_{\Sigma_c} &= (2.45 \sim 2.94) \text{ GeV} , \\ M_{\Sigma_b} &= (5.70 \sim 6.62) \text{ GeV} , \\ M_{\Sigma_c} - M_{\Lambda_c} &\leq 207 \text{ MeV} , \\ M_{\Sigma_b} - M_{\Lambda_b} &\leq 163 \text{ MeV} , \end{aligned} \quad (25)$$

which we confirm. However, we have not tried to improve these results for the absolute values of the masses due to the ambiguity in the definition of the heavy quark mass input mentioned earlier at LO, which induces large errors.

$$\Xi_c(cs q) / \Lambda_c(cqq)$$

– *Optimal choice of the currents:* we start from the general choice of interpolating currents given in Eq. (1). We study the b -behaviour of the predictions in Fig. 1a), by fixing $\tau = 0.35 \text{ GeV}^{-2}$ and $t_c = 15 \text{ GeV}^2$. The result presents b -stability around $b = 0$, which is in the range given in Eq. (3) obtained from light baryons and heavy non-strange baryons. We consider this value as the optimal choice of the interpolating currents. However, this generous range does not favour the *ad hoc* choice around




Figure 1. Ξ_c/Λ_c : a) b -behaviour of the double ratio of sum rules (DR) given $\tau = 0.35 \text{ GeV}^{-2}$ and $t_c = 15 \text{ GeV}^2$: r_1^{ds} dashed-dotted (blue), r_2^{sd} dotted (green), r_{21}^{sd} continuous (red); b) τ -behaviour for $b = 0$ and $t_c = 15 \text{ GeV}^2$; c) t_c -behaviour of the DR given $b = 0$ and $\tau = 0.35 \text{ GeV}^{-2}$; We have used $\kappa = 0.7$.

1 used in the existing literature [14,15]. Therefore, in this channel, we shall work with:

$$b \simeq 0. \quad (26)$$

– τ stabilities: we show in Fig. 1b) the τ behaviour of the different DR at fixed $t_c = 15 \text{ GeV}^{-2}$ and $b = 0$.

– t_c stabilities and choice of the sum rules: fixing $b = 0$ from the previous analysis, we study in Fig. 1c) the t_c -behaviour of predictions. Among the three sum rules, we retain r_{21}^{sd} which is the most stable in t_c and then less affected by the higher state contributions.

– *Results:* From this r_{21}^{sd} sum rule, we can deduce the DR:

$$r_{\Xi_c}^{sd} = 1.080(10)(2)(6)(2)(1.5), \quad (27)$$

where the value $\kappa = 0.7$ has been taken. We have considered the mean value of r_{21}^{sd} from $t_c = 10$ to 20 GeV^2 . The errors are due respectively to the values of t_c , $\tau = (0.35 \pm 0.05) \text{ GeV}^{-2}$, m_c , m_s and the factorization of the four-quark condensate ρ . The errors due to b and some other SU(3) symmetric QCD parameters are negligible. The large error due to κ compiled in Table 1 is not included in Eqs. (27). Using as input the data [9]:

$$M_{\Lambda_c}^{exp} = (2286.46 \pm 0.14) \text{ MeV}, \quad (28)$$

and adding the different errors quadratically, one can deduce:

$$M_{\Xi_c} = (2469.4 \pm 26.6) \text{ MeV}, \quad (29)$$

which agrees nicely with the data [9]:

$$M_{\Xi_c}^{exp} = (2467.9 \pm 0.4) \text{ MeV} . \quad (30)$$

For improving the existing value of κ , we allow a 1σ deviation of the DR prediction from the experimental value. In this way, we deduce:

$$\kappa = 0.700(50) . \quad (31)$$

$$\Xi_b(bsq)/\Lambda_b(bqq)$$

We repeat the previous analysis in the case of the b -quark. The analysis of the ratio of sum rules shows similar curves than for the charm case except the obvious change of scale. Using r_{21}^{sd} , we illustrate the analysis using $\kappa = 0.74$. In this way, we obtain:

$$r_{\Xi_b}^{sd} = 1.030(2.5)(0.5)(1.5)(0.5)(0.5) , \quad (32)$$

where the errors come from t_c from 45 to 80 GeV^2 , $\tau = (0.18 \pm 0.05) \text{ GeV}^{-2}$, m_b , m_s and the factorization of the four-quark condensate ρ . From this result, and using:

$$M_{\Lambda_b}^{exp} = (5620.2 \pm 1.6) \text{ MeV} , \quad (33)$$

we deduce:

$$M_{\Xi_b} = (5789 \pm 16) \text{ MeV} , \quad (34)$$

which agrees quite well with the data [9]:

$$M_{\Xi_b}^{exp} = (5792.4 \pm 3.0) \text{ MeV} . \quad (35)$$

Allowing a 1σ deviation from the data, we deduce:

$$\kappa = 0.738(23) . \quad (36)$$

$$\Omega_c(css)/\Sigma_c(cqq)$$

We do an analysis similar to the one in the previous section. The result for the c -quark is shown in Fig 2. One can notice that the optimal choice of the current is the same as in Eq. (2) which we fix to the value $b=0$. One can notice from Fig 2a) and Fig 2b), that only r_2^{sd} presents simultaneously τ and b stabilities from which we extract the optimal result. Using $\kappa = 0.78$, the final result from r_2^{sd} is:

$$r_{\Omega_c}^{sd} = 1.111(1.4)(1.3)(16.4)(0.2) . \quad (37)$$

The errors are due respectively to the values of $\tau = (1.0 \sim 1.2) \text{ GeV}^{-2}$, m_c , m_s and the factorization of the four-quark condensate ρ . The one due to t_c from 10 to 20 GeV^2 is negligible. Using this previous result together with the experimental averaged value [9]:

$$M_{\Sigma_c}^{exp} = (2453.6 \pm 0.25) \text{ MeV} , \quad (38)$$

one can deduce:

$$M_{\Omega_c} = 2726.9(40.5) \text{ MeV} , \quad (39)$$

in good agreement with the data:

$$M_{\Omega_c}^{exp} = (2697.5 \pm 2.6) \text{ MeV} . \quad (40)$$

Now, we study the influence of κ on the mass prediction. Allowing a 1σ deviation from the experimental mass, we deduce the estimate:

$$\kappa = 0.775(15) . \quad (41)$$




Figure 2. Ω_c/Σ_c : a) τ -behaviour of DR given $b=0$ and $t_c=14 \text{ GeV}^2$: r_1^{ds} dashed-dotted (blue), r_2^{ds} dotted (green), r_{21}^{ds} continuous (red); b) b -behaviour of the DR given $\tau=0.6 \text{ GeV}^{-2}$ (τ -stability of r_1^{ds}) and $t_c=14 \text{ GeV}^2$; c) t_c -behaviour of r_2^{ds} given $b=0$ and $\tau=1 \text{ GeV}^{-2}$.

Final value of κ

Taking the (arithmetic) mean value of $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$ from the different channels $\Xi_{c,b}$ [Eqs. (31) and (36)] and Ω_c [Eq. (41)], we deduce:

$$\kappa = 0.738(29) , \quad (42)$$

which we can consider as an improved estimate of this quantity compared with the existing one from the light mesons [31,1] and baryons [20]:

$$\kappa = 0.700(100) . \quad (43)$$

6. The mass of the $\Omega_b(bss)$

We repeat the previous analysis of the $\Omega_c(css)$ in the case of the b -quark. The curves present the same qualitative behaviour as in the case of the charm, where, only r_2^{ds} survives the different tests of stabilities. The optimal value is taken at the extremum $\tau = (0.25 \pm 0.05) \text{ GeV}^{-2}$ and in the t_c stability region. Then, we obtain:

$$r_{\Omega_b}^{sd} = 1.0455(20)(22)(41)(13)(37) . \quad (44)$$

The errors are due respectively to the values of τ , m_b , m_s and the factorization of the four-quark condensate ρ . The last error is due to κ . The one due to t_c in the t_c stability region is negligible. Using this value together with the experimental averaged value [9]:

$$M_{\Sigma_b}^{exp} = 5811.2 \text{ MeV} , \quad (45)$$

one can deduce the result:

$$M_{\Omega_b} = 6075.6(37.2) \text{ MeV} , \quad (46)$$




Figure 3. Ω_b/Σ_b : a) τ -behaviour of DR given $b = 0$ and $t_c = 60$ GeV^2 : r_1^{ds} dashed-dotted (blue), r_2^{ds} dotted (green), r_{21}^{ds} continuous (red); b) b -behaviour of the DR given $\tau = 0.25$ GeV^{-2} (τ -stability of r_1^{ds} and $t_c = 60$ GeV^2); c) t_c -behaviour of r_2^{ds} given $b = 0$ and $\tau = 0.25$ GeV^{-2} . We have used $\kappa = 0.738$.

which we compile in Table 2. This result agrees within the errors with the one from the CDF collaboration [37]: 6054.4(6.9) MeV but disagrees by about 2.4σ with the D0 value [38]: $M_{\Omega_b}^{D0} = 6165.0(13.0)$ MeV given in the same table.

7. The mass of the $\Xi'_{c,b}(Qsq)$

We do a similar analysis for the $\Xi'_{c,b}$, which we show in Fig. 4, where we have only retained the r_2^{ds} which satisfies all stability tests. We obtain:

$$r_{\Xi'_c}^{sd} = 1.043(1)(2)(6)(2)(3)(7), \quad (47)$$

The errors are due respectively to the values of $\tau = (0.9 \pm 0.1)$ GeV^{-2} , m_c , m_s , ρ , $b = -(0.4 \pm 0.2)$, and κ . Using the experimental value of M_{Σ_c} , we obtain:

$$M_{\Xi'_c} = 2559(25) \text{ MeV}, \quad (48)$$

which we compile in Table 2. Our prediction is in good agreement (1σ) with the data [9]:

$$M_{\Xi'_c}^{exp} = 2576(3.1) \text{ MeV}. \quad (49)$$

A similar analysis is done for Ξ'_b , which is summarized in Fig. 5. One can notice that here the b -stability occurs at 0 in this channel and there is no sharp selection between r_1^{ds} and r_2^{ds} . We obtain the mean from r_1^{ds} and r_2^{ds} :

$$r_{\Xi'_b}^{sd} = 1.014(3.4)(5)(1.7)(0.5)(2)(0.5)(3), \quad (50)$$

where the sources of the errors are $\tau = (0.5 \pm 0.1)$ GeV^{-2} , m_b , m_s , ρ , $b = (0 \pm 0.2)$, and κ . The last




Figure 4. Ξ'_c/Σ_c : a) τ -behaviour of DR given $b = -0.4$ and $t_c = 14$ GeV^2 : r_1^{ds} dashed-dotted (blue), r_2^{ds} dotted (green), r_{21}^{ds} continuous (red); b) b -behaviour of the DR given $\tau = 0.9$ GeV^{-2} and $t_c = 14$ GeV^2 ; c) t_c -behaviour of r_2^{ds} given $b = -0.4$ and $\tau = 1$ GeV^{-2} . We have used $\kappa = 0.74$.

error comes from the choice of the sum rules. Using the experimental value of M_{Σ_b} , we predict:

$$M_{\Xi'_b} = 5893(42) \text{ MeV}, \quad (51)$$

which we compile in Table 2.

8. The masses of the spin 3/2 baryons

As a preliminary step of the analysis, we check the different results obtained in [5]:

$$\begin{aligned} M_{\Sigma_c^*} &= (2.15 \sim 2.92) \text{ GeV}, \\ M_{\Sigma_b^*} - M_{\Sigma_c^*} &= 3.3 \text{ GeV}, \end{aligned} \quad (52)$$

and confirm them. However, like in the spin 1/2 case, we have not tried to improve these (old) results.

$$\Xi_c^*(csq)/\Sigma_c^*(cqq)$$

We repeat the previous DR analysis for the case of the Ξ_c^* . We show in Fig. 6a) the τ -behaviour of the mass predictions for $t_c = 14$ GeV^2 . From this analysis, we do not retain r_{21}^{ds} which differs completely from r_1^{ds} and r_2^{ds} , while we do not consider r_1^{ds} which is τ -unstable. We show in Fig. 6b) the t_c -behaviour of r_2^{ds} given $\tau = 0.9$ GeV^{-2} .

We deduce the optimal value:

$$r_{\Xi_c^*}^{sd} = 1.049(1)(10)(4)(4)(17.5)(1). \quad (53)$$

The errors are due respectively to the values of $\tau = (0.9 \pm 0.1)$ GeV^{-2} , m_c , m_s , ρ (factorization of the four-quark condensate) and $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle = 0.74 \pm 0.03$. The




Figure 5. Ξ_b^*/Σ_b^* : a) τ -behaviour of DR given $b = 0$ and $t_c = 60$ GeV^2 : r_1^{ds} dashed-dotted (blue), r_2^{ds} dotted (green), r_{21}^{ds} continuous (red); b) b -behaviour of the DR given $\tau = 0.5$ GeV^{-2} and $t_c = 60$ GeV^2 ; c) t_c -behaviour of r_2^{ds} given $b = 0$ and $\tau = 0.5$ GeV^{-2} . We have used $\kappa = 0.74$.

ones due to some other parameters are negligible. Using the data [9]:

$$M_{\Sigma_c^{exp}} = (2517.97 \pm 1.17) \text{ MeV} , \quad (54)$$

and adding the different errors quadratically, we deduce the results in Table 2.

$$\Xi_b^*(bsq)/\Sigma_b^*(bqq)$$

We extend the analysis to the case of the bottom quark. The curves are qualitatively similar to the charm case. We deduce:

$$r_{\Xi_b}^{sd} = 1.022(2)(2)(0.5)(1)(2) . \quad (55)$$

The sources of the errors are the same as for the Ξ_c^* , where here $\tau = (0.25 \pm 0.05) \text{ GeV}^{-2}$ and m_c replaced by m_b . The ones due to some other parameters are negligible. Using the averaged data [9]:

$$M_{\Sigma_b^{exp}} = (5832.7 \pm 6.5) \text{ MeV} , \quad (56)$$

and adding the different errors quadratically, we deduce:

$$M_{\Xi_b^*} = (5961 \pm 21) \text{ MeV} , \quad (57)$$

which we report in Table 2.

$$\Omega_c^*(css)/\Sigma_c^*(cqq)$$

We pursue the analysis to the case of the $\Omega_c^*(css)$. We show the τ -behaviour of the different DR in Fig. 7a). From this figure, we shall not retain r_{21}^{ds} which differs




Figure 6. Ξ_c^*/Σ_c^* : a) τ -behaviour of the double ratio of sum rules (DR) by giving $t_c = 14$ GeV^2 : r_1^{ds} dashed-dotted (blue), r_2^{sd} dotted (green) and r_{21}^{sd} continuous (red); c) t_c -behaviour of r_2^{ds} for a given optimal $\tau = 0.9$ GeV^{-2} .

completely from r_1^{ds} and r_2^{ds} . Given the optimal value of $\tau = 1$ GeV^{-2} , we show the t_c -behaviour of the DR r_1^{ds} and r_2^{ds} in Fig. 7b). The final result is the mean from r_1^{ds} and r_2^{ds} :

$$r_{\Omega_c}^{sd} = 1.109(3)(10)(10)(4)(0.5)(8) . \quad (58)$$

The 1st error is due to the choice of r_i^{sd} . The other ones are due to $\tau = (1.0 \pm 0.2) \text{ GeV}^{-2}$, m_c , m_s , ρ and κ . The other QCD parameters give negligible errors. Using the averaged data in Eq. (56), and adding the different errors quadratically, one can deduce the result in Table 2.

$$\Omega_b^*(bss)/\Sigma_b^*(bqq)$$

We repeat the previous analysis in the b -channel. The curves are qualitatively analogue to the ones of the charm. We shall not consider r_{21}^{sd} because of its incompatibility with the other ones. From the mean of r_1^{ds} and r_2^{ds} , we deduce:

$$r_{\Omega_b}^{sd} = 1.040(4)(2)(4.6)(0.2)(6) , \quad (59)$$

where the sources of the errors are the same as for Ω_c^* , where $\tau = (0.30 \pm 0.05) \text{ GeV}^{-2}$ here and m_c replaced by m_b . Using the averaged data in Eq. (56), and adding the different errors quadratically, one can deduce:

$$M_{\Omega_b^*} = (6066 \pm 49) \text{ MeV} , \quad (60)$$

which we report in Table 2.

9. Hyperfine mass-spillings

Combining the results for spin 1/2 and spin 3/2 given in Table 2, we deduce in Table 3 the values of the hyperfine mass-spillings. From our analysis, one expects that the Ω_Q^* can only decay electromagnetically to $\Omega_Q + \gamma$ due to phase space, while the Ξ_Q^* can, in addition, decay hadronically to $\Xi_Q + \pi$. On one hand, our result for the




Figure 7. Ω_c^*/Σ_c^* : a) τ -behaviour of the double ratio of sum rules (DR) by giving $t_c = 14 \text{ GeV}^2$: r_1^{ds} dashed-dotted (blue), r_2^{sd} dotted (green) and r_{21}^{sd} continuous (red); b) t_c -behaviour of the DR r_1^{ds} and r_2^{sd} giving $\tau = 1 \text{ GeV}^{-2}$.

Table 2

QSSR predictions of the strange heavy baryon masses in units of MeV from the double ratio (DR) of sum rules with the QCD input parameters in Table 1 and using as input the observed masses of the associated non-strange heavy baryons. We have used $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle = 0.74 \pm 0.03$ fixed from the experimental $\Xi_{c,b}$ and Ω_c masses.

Baryons	I	$r_{B^*Q}^{sd}$	Mass	Data
$J^P = \frac{1}{2}^+$				
$\Xi_c(cqs)$	$\frac{1}{2}$		input	2467.9 ± 0.4 [PDG]
$\Omega_c(css)$	0		input	2697.5 ± 2.6 [PDG]
$\Xi_b(bqs)$	$\frac{1}{2}$		input	5792.4 ± 3.0 [PDG]
$\Xi'_c(cqs)$	$\frac{1}{2}$	1.043(10)	2559(25)	2575.7 ± 3.1 [PDG]
$\Xi'_b(bqs)$	$\frac{1}{2}$	1.014(7)	5893(42)	—
$\Omega_b(bss)$	0	1.0455(64)	6076(37)	6165.0 ± 13 [D0] 6054.4 ± 6.9 [CDF]
$J^P = \frac{3}{2}^+$				
$\Xi_c^*(cqs)$	$\frac{1}{2}$	1.049(8)	2641(21)	2646.1 ± 1.3 [PDG]
$\Omega_c^*(css)$	0	1.109(17)	2792(38)	2768.3 ± 3.0 [PDG]
$\Xi_b^*(bqs)$	$\frac{1}{2}$	1.024(8)	5961(21)	—
$\Omega_b^*(bss)$	0	1.040(9)	6066(49)	—

unobserved mass-differences $M_{\Omega_b^*} - M_{\Omega_b}$, $M_{\Xi_b^*} - M_{\Xi_b}$ agree within the errors with the ones from quark models [11] and $1/N_c$ expansion [19] and seems to behave like $1/m_b$ despite the large errors. On the other hand, we predict with a better precision $M_{\Xi_b^*} - M_{\Xi_b} \simeq M_{\Xi_c^*} - M_{\Xi_c}$. A future precise measurement of the Ξ'_b , Ξ_b^* and Ω_b^* will shed light on the quark mass behaviour of these mass-differences, which we also plan to study in a future work.

10. Summary and Conclusions

We have directly extracted (for the first time) the heavy baryons (charmed $C = 1$ and bottom $B = -1$) mass-splittings due to $SU(3)$ breaking using double ratios (DR) of QCD spectral sum rules(QSSR), which are less sensitive to the exact value and the definition of the

Table 3

QSSR predictions of the strange heavy baryon hyperfine splittings in units of MeV from the double ratio (DR) of sum rules with the QCD input parameters in Table 1 and using as input the predicted values in Table 2. We have added the errors quadratically.

Hyperfine Splittings	Data
$M_{\Xi_c^*} - M_{\Xi_c} = 173(21)$	179(1)
$M_{\Xi_c^*} - M_{\Xi_c'} = 82(33)$	70(3)
$M_{\Omega_c^*} - M_{\Omega_c} = 95(38)$	70(3)
$M_{\Xi_b^*} - M_{\Xi_b} = 169(21)$	—
$M_{\Xi_b^*} - M_{\Xi_b'} = 68(47)$	—
$M_{\Omega_b^*} - M_{\Omega_b} = -10(61)$	$M_{\Sigma_b^*} - M_{\Sigma_b} = 22$

heavy quark mass, to the perturbative radiative corrections and to the QCD continuum contributions than the simple ratios commonly used in the current literature for determining the heavy baryon masses:

Remarking that the leading term controlling the mass-splittings is, in most of the cases, the ratio $\kappa \equiv \langle \bar{s}s \rangle / \langle \bar{d}d \rangle$ of the condensate rather than the running mass $\bar{m}_s$, we use as input the observed masses of the $\Xi_{c,b}$ and Ω_c , for extracting κ . We obtain the mean value from Eqs. (31), (36) and (41):

$$\kappa = 0.738(29) \text{ [Eq. (42)]}, \quad (61)$$

which we can consider as an improved estimate of this quantity compared with the existing one $\kappa = 0.7 \pm 0.1$ compiled in Table 1 from the light mesons [31,1] and baryons [20].

Using this value of κ , we give predictions of the $\Xi'_{c,b}$, Ω_b and spin 3/2 baryons masses which are summarized in Table 2. These predictions are in good agreement with the experimental masses and can be considered as improvements of existing QSSR results based on the simple ratio of moments [14,15].

Our result for the Ω_b favours the one observed by CDF [37] but disagrees within 2.4σ with the one from D0 [38]. Most of predictions agree with the ones from different approaches (quark models [11], lattice calculations [18] and large N_c [19]). Our predictions for the not yet observed states Ξ'_b , Ξ_b^* and Ω_b^* given in Table 2 can serve in a near future as a test of the QSSR approach.

We show in Table 3 our predictions for the hyperfine splittings. Our results agree with the observed values, while the ones for not yet observed states can serve as a test of the QSSR approach. From our results, we expect that the Ω_Q^* can only decay electromagnetically to $\Omega_Q + \gamma$ due to the available phase space, while the Ξ_Q^* can, in addition, decay hadronically to $\Xi_Q + \pi$.

One can also notice that, if we have used a $SU(3)$ symmetric quark condensates $\langle \bar{s}s \rangle \simeq \langle \bar{d}d \rangle$, the predictions would be systematically lower by about (70 ~ 100) MeV than the predictions given in Table 2. In this case, the agreement with the observed masses in different channels cannot be achieved. It would be interesting to understand analogous effects of κ using some other approaches.

Acknowledgements

R.M.A and M.N. acknowledge the financial support from FAPESP and CNPq (Brazil). M.N. also thanks

S. Narison for the hospitality at the Laboratoire de Physique Théorique et Astroparticules of Montpellier. S.N. has been partially supported by the CNRS-IN2P3 within the project “Hadron Physics in QCD” and within the French-China Particle Physics Laboratory (FCPPL). We thank Jean-Marc Richard for some discussions.

REFERENCES

- For reviews, see e.g.: S. Narison, *Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol.* **17** (2004) 1-778 [hep-ph/0205006]; S. Narison, *World Sci. Lect. Notes Phys.* **26** (1989) 1 ; S. Narison, *Acta Phys. Pol.* **B 26** (1995) 687; S. Narison, *Riv. Nuovo Cim.* **10 (N2)** (1987) 1; S. Narison, *Phys. Rept.* **84** (1982) 263 .
- J.S. Bell and R.A. Bertlmann, *Nucl. Phys.* **B 177** (1981) 218. **B187** (1981) 285; R.A. Bertlmann, *Acta Phys. Austr.* **53** (1981) 305.
- M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B 147** 385, 448 (1979).
- E. Bagan, M. Chabab, H.G. Dosch, S. Narison, *Phys. Lett.* **B 278** (1992) 367.
- E. Bagan, M. Chabab, H.G. Dosch, S. Narison, *Phys. Lett.* **B 287** (1992) 176.
- E. Bagan, M. Chabab, S. Narison, *Phys. Lett.* **B 306** (1993) 350.
- E. Bagan, M. Chabab, H.G. Dosch, S. Narison, *Phys. Lett.* **B 301** (1993) 243.
- The D0 collaboration: V. Abazov et al. *Phys. Rev. Lett.* **99** (2007) 052001; *arXiv: 0808.4142 [hep.ex]*; The CDF collaboration: D. Acosta et al., *Phys. Rev. Lett.* **96** (2006) 202001; T. Aaltonen et al. *Phys. Rev. Lett.* **99** (2007) 052002; *ibid. Phys. Rev. Lett.* **99** (2007) 202001.
- PDG, C. Amsler et al., *Phys. Lett.* **B 667** (2008) 1.
- The Babar collaboration: B. Aubert et al., *Phys. Rev. Lett.* **97** (2006) 232001; *ibid Phys. Rev. Lett.* **99** (2007) 202001; The Belle collaboration: R. Chistov et al., *Phys. Rev. Lett.* **97** (2006) 162001.
- For recent reviews, see e.g.: M. Karliner, B. Keren-Zur, J.H. Lipkin, J.L. Rosner, *Annals Phys.* **324** (2009) 2; E. Klempt, J.M. Richard, *arXiv:0901.2055 [hep-ph]* and references therein.
- R.S. Marques de Carvalho et al., *Phys.Rev.* **D 60** (1999) 034009.
- F. Hussain, J.G. Korner, M. Kramer, G. Thompson, *Z. Phys.* **C 51** (1991) 321.
- F.O. Durães, M. Nielsen, *Phys. Lett.* **B 658** (2007) 40.
- X. Liu, H-X. Chen, Y-R. Liu, A. Hosaka, S-L. Zhu, *arXiv:0710.0123 [hep-ph]*; Z-G. Wang, *arXiv:0809.3038 [hep-ph]*; *ibid, arXiv:0704.1106 [hep-ph]*; J-R. Zhang, M-Q. Huang, *arXiv:0811.3266 [hep-ph]*, *ibid, arXiv:0810.5396 [hep-ph]*; *ibid, arXiv:0805.0479 [hep-ph]*.
- S. Groote, J.G. Korner, O.I. Yakovlev, *Eur. Phys. J.* **C 58** (2008) 355.
- D. Ebert, R.N. Faustov and V.O. Galkin, *Phys. Rev.* **D 72** (2005) 034026.
- W. Detmold, C-J. D. Lin, M. Wingate, *arXiv:0812.2583 [hep-lat]*; R. Lewis, R.M. Woloshyn, *Phys. Rev.* **D 79** (2009) 014502; H. Na, S. Gottlieb, *arXiv:0812.1235 [hep-lat]*.
- E. Jenkins, *Phys. Rev.* **D 77** (2008) 034012.
- Y. Chung et al. *Z. Phys.* **C 25** (1984) 151; H.G. Dosch, M. Jamin and S. Narison, *Phys. Lett.* **B 220** (1989) 251.
- B.L. Ioffe, *Nucl. Phys.* **B 188** (1981) 317 , **B 191** (1981) 591; A.A.Ovchinnikov and A.A.Pivovarov, *Yad. Fiz.* **48** (1988) 1135.
- G. Launer, S. Narison and R. Tarrach, *Z. Phys.* **C 26** (1984) 433.
- S. Narison, *Phys. Lett.* **B 210** (1988) 238; *Phys. Lett.* **B 605** (2005) 319.
- S. Narison, *Phys. Lett.* **B 322** (1994) 327.
- S. Narison, *Phys. Lett.* **B 337** (1994) 166; *Phys. Lett.* **B 668** (2008) 308; *Phys. Lett.* **B 327** (1994) 354.
- S. Narison, *Phys. Lett.* **B 358** (1995) 113; *Phys. Lett.* **B 466** (1999) 345.
- E.G. Floratos, S. Narison and E. de Rafael, *Nucl. Phys.* **B 155** (1979) 155.
- S. Narison and V.I. Zakharov, *Phys. Lett.* **B 679** (2009) 355; K. Chetyrkin, S. Narison and V.I. Zakharov, *Nucl. Phys.* **B 550**, 353 (1999); For a review, see e.g. S. Narison, *Nucl. Phys. Proc. Suppl.* **164** (2007) 225.
- For a review, see e.g. V.A. Zakharov, *Nucl. Phys. (Proc. Sup.)* **B 74** (1999) 392; V.I. Zakharov, *Nucl. Phys. Proc. Suppl.* **164** (2007) 240; V.I. Zakharov, contribution at QCD 08 (Montpellier); R. Akhoury and V.I. Zakharov, *Phys. Lett.* **B 438** (1998) 165; F.V. Gubarev, M.I. Polikarpov and V.I. Zakharov, hep-th/9812030.
- S. Narison, *Phys. Lett.* **B 673** (2009) 30.
- For reviews, see e.g.: S. Narison, *Phys.Rev.* **D 74** (2006) 034013; *arXiv:hep-ph 0202200*; *Phys. Lett.* **B 216** (1989) 191; *Phys. Lett.* **B 358** (1995) 113; *Phys. Lett.* **B 466** (1999) 345; S. Narison, H.G. Dosch, *Phys. Lett.* **B 417** (1998) 173; S. Narison, N. Paver, E. de Rafael and D. Treleani, *Nucl. Phys.* **B 212** (1983) 365; S. Narison, E. de Rafael, *Phys. Lett.* **B 103** (1981) 57; C. Becchi, S. Narison, E. de Rafael, F.J. Yndurain, *Z. Phys.* **C 8** (1981) 335.
- S. Narison, *Phys. Lett.* **B 300** (1993) 293; *ibid B 361* (1995) 121.
- R.A. Bertlmann, G. Launer and E. de Rafael, *Nucl. Phys.* **B 250** (1985) 61; R.A. Bertlmann et al., *Z. Phys.* **C 39** (1988) 231.
- F.J. Yndurain, hep-ph/9903457.
- S. Narison, *Phys. Lett.* **B 387** (1996) 162.
- S. Narison, *Phys. Lett.* **B 197** (1987) 405; *Phys. Lett.* **B 341** (1994) 73; *Phys. Lett.* **B 520** (2001) 115.
- The CDF collaboration: T. Aaltonen et al., *arXiv: 0905.3123 [hep.ex]*.
- The D0 collaboration: V.M. Abazov et al., *arXiv: 0808.4142 [hep.ex]*